

Vision and Comfort over 16 hours of wear of a Water Surface Treated Soft Contact Lens for Astigmatism

Jennifer Fogt, OD MS Abigail Menner OD Kate McClure OD MS Kimberly Patton OD MS




This study enrolled current wearers of soft toric contact lenses to assess their wear experience when wearing a daily disposable contact lens with a water surface treatment over 16 hours of wear time.

METHODS

35 participants with astigmatism in at least one eye were consented and completed the study. The initial visit evaluated ocular heath and visual acuity to ensure inclusion/exclusion criteria were met. Those eligible were refit into the study lenses (PRECISION1® for Astigmatism, Alcon, Fort Worth TX).

After 1 week of lens wear, participants returned for a follow-up visit. After a successful follow-up visit, participants were scheduled to complete surveys throughout the evening of 5 weekdays.



Participants used personal smart devices to enter the time of insertion of their lenses each morning and then responded to surveys of their comfort and vision while wearing the study lenses at 10, 12, 14 and 16 hours of lens wear.

Evening surveys included 1-10 forced choice grading of comfort and quality of vision, with 1 labeled as "poor" and 10 labeled as "excellent." Evening scores for each time point on the five days of surveys were averaged for each participant and overall median and interquartile range was calculated for each study time point.

RESULTS

Thirty participants were fit with the study lenses for astigmatism in both eyes.

Mean (standard deviation) LogMAR acuity OU with the study lenses was -0.19, which equates to 20/12.5⁻¹ Snellen acuity.

Visual Acuity with study lenses								
n=30	OD	OS	OU					
Mean LogMAR acuity	-0.12	-0.14	-0.19					
Standard deviation	0.08	0.08	0.06					
Mean Snellen equivalent	20/16+1	20/16+2	20/12.5 ⁻¹					

Rotation of lenses was less than 10 degrees in 56/61 eyes (91.8%) of assessed.

For the evening survey (n=34), the median score (interquartile range) for comfort was 9.5 (1.6) 10 hours; 9.2 (1.8) at 12 hours; 8.9 (1.7) at 14 hours; and 8.3 (2.6) at 16 hours of wear. Median quality of vision scores were 9.9 (1.2) at 10 hours; 9.6 (1.6) at 12 hours; 9.3 (1.6) at 14 hours; and 8.9 (2.6) at 16 hours of wear.

Evening Surveys										
	10 hour Comfort Score	10 hour Vision score	12 hour Comfort Score	12 hour Vision score	14 hour Comfort Score	14 hour Vision score	16 hour Comfort Score	16 hour Vision score		
Median score	9.5	9.9	9.2	9.6	8.9	9.3	8.3	8.9		
Interquartile range	1.6	1.2	1.8	1.6	1.7	1.6	2.6	2.6		

CONCLUSIONS

In this study, all participants with astigmatism had visual acuity of 20/20 or better when wearing the daily disposable study lenses with water surface treatment.

Participants had median subjective comfort scores greater than 8 and quality of vision scores greater than 9 on a scale of 1-10 at 16 hours of lens wear.

ACKNOWLEDGEMENTS

Funding for this research was provided by Alcon. The investigators designed and implemented this study.

The project described was supported in part by Award Number Grant UL1TR002733 from the National Center for Advancing Translational Sciences. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Center for Advancing Translational Sciences or the National Institutes of Health.

BIBLIOGRAPHY

- 1. Sulley A, Young G, Hunt C. Factors in the success of new contact lens wearers. *Contact Lens Anterio*. 2017;40(1):15-24.
- 2. Nichols JJ, Willcox MD, Bron AJ, et al. The TFOS International Workshop on Contact Lens Discomfort: executive summary. *Invest Ophthalmol Vis Sci.* 2013;54(11):TFOS7-TFOS13.
- 3. Dumbleton K, Woods CA, Jones LW, Fonn D. The impact of contemporary contact lenses on contact lens discontinuation. *Eye Contact Lens.* 2013;39(1):93-99.
- 4. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)-A metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform.* 2009;42(2):377-381.