



Javier Rojas¹, David Piñero², Javier Sebastián³, Julio Ezpeleta⁴, Alejandra Consejo⁵

Introduction

Purpose

Several authors have attempted to measure the CSJ angle using optical coherence tomography (OCT) [1-3]. Seguí-Crespo et al. and Hall et al. measured the angle by means of a point-and-click calliper that an observer manually manipulated to locate the CSJ and measure the angle [1,2]. Tan et al. went a step further and developed an algorithm to automatically measure the angle, although the CSJ was still located manually by an observer [3]. They described good inter-observers [3]. Moreover, measurements of the CSJ angle in these studies were taken at a single point of each quadrant.

The aim of this study was to introduce a fully objective, automated methodology to estimate CSJ angle in 360 degrees in the limbal position, assessed from 3-dimensional corneoscleral topography [4,5]. This methodology was used to evaluate the mean CSJ angle in healthy eyes objectively.

17.8°).

The corneoscleral topography of 105 healthy right eyes of Caucasian subjects (67% women and 33% men) aged between 18 and 59 years were retrospectively analysed. These eyes were previously measured with the Eye Surface Profiler (ESP, Eaglet Eye, The Netherlands). The raw anterior eye height data (x, y, and z coordinates) were exported to build three-dimensional corneoscleral topography maps in a four-step process:

- 1.The limbus position was calculated in 360 semi-meridians using a purpose-designed algorithm [4,5].
- 2.After limbus demarcation, auxiliary points were placed 0.6 mm horizontally away from the limbus (yellow squares in Figure 1). Angle α (see Figure 1) was evaluated as the arctangent of the adjacent, i.e., 0.6 mm, and the opposite a, calculated as the distance between the corresponding auxiliary points. The same procedure was repeated to estimate angle β (see Figure 1).
- 3. In the following step, angle ϕ (see Figure 1) was calculated as $\phi = 180^\circ - \alpha$ (see Figure 1).
- 4. Finally, the CSJ angle was obtained: $CSJ = \phi + \beta$ (see Figure 1).

Discussion and conclusions

References

The CSJ angle influences sagittal height, a key parameter lenses. However, to date, only a few works have characterised this parameter [2,6]. These are based on manually positioning virtual callipers on an image, which makes the process subjective and poorly repeatable [7]. This novel method is designed to obtain a more complete, realistic description of the transition from the cornea to the sclera than the current standards, and to avoid the loss of accuracy inherent in subjective criteria [8]

- Int Ophthalmol 2019;39:2603-2609.

Corneoscleral junction angle in healthy eyes assessed objectively

Methods



Figure 1. Methodology for corneoscleral junction (CSJ) angle calculation. The solid black line corresponds to the corneoscleral profile in one out of 360 semi-meridians. For details on angle estimation see the text.

1. Seguí-Crespo M, Ariza-Gracia MA, Sixpene NL, Piñero DP. Geometrical characterization of the corneo-scleral transition in normal patients with Fourier domain optical coherence tomography.

2. Hall LA, Hunt C, Young G, Wolffsohn J. Factors affecting corneoscleral topography. Invest Ophthalmol Vis Sci 2013;54:3691–3701.

3. Tan B, Graham AD, Tsechpenakis G, Lin MC (2014) A novel analytical method using OCT to describe the corneoscleral junction. Optom Vis Sci 91:650–657.

4. Consejo A. Llorens-Quintana C. Radhakrishnan H, Iskander RD. Mean shape of the human limbus. J Cataract Refract Surg. 2017;43(5):667-672.

5. Consejo A, Iskander RD. Corneo-scleral limbus demarcation from 3D height data. Cont Lens Anterior Eye. 2016;39(6):450-457.

6. Hall LA, Young G, Wolffsohn JS, et al. The influence of corneoscleral topography on soft contact lens fit. Invest Ophthalmol Vis Sci 2011;52:6801–6806

7. Bergmann B, Wolffsohn JS, Bandlitz S. Scheimpflug imaging for grading and measurement of corneo-scleral-profile in different quadrants. Contact Lens and Anterior Eye. 2022 Aug 24:101753. 8. Baumeister M, Terzi E, Ekici Y, Kohnen Y. Comparison of manual and automated methods to determine horizontal corneal diameter. J Cataract Refract Surg. 2004;30:374-38

Results Quadra ✓The group's mean CSJ angle was 177.5 ± **1.1°**, Nasa ✓Regional differences were observed Tempo (Table 1) Superi ✓The CSJ angle was rotationally Inferio asymmetric (Figure 2). There was a mean 7.7 \pm 3.7° difference between the steepest (smallest) and flattest (largest) angle within the same eye (greatly depended on 185

✓The CSJ angle was smaller (steeper) in the nasal region than in the remaining sectors.

the individual as it ranged from 3.5° to

✓The CSJ angle and limbal radius provided by the ESP were moderately correlated (r=0.43, p<0.001).





¹GOO MSc, FAAO, Centro de Lentes de Contacto - Natural Optics Balaguer (Spain) ²PhD, Universidad de Alicante (Spain) ³GOO, MSc, QVisión - Óptica Ronda (Spain) ⁴DOO, Óptica Julio Ezpeleta (Spain) ⁵PhD, Universidad de Zaragoza (Spain)

Email contact: <u>soyjavirojas@hotmail.com</u>, <u>alejandra.consejo@unizar.es</u> Disclosures: Javier Rojas, consultant of Eaglet Eye



Universidad Zaragoza

ant	Mean CSJ angle ± SD (°)	Range (°)	p-value (paired t-test)
l	176.4 ± 1.1	[172.9, 178.7]	<0.001
ral	178.2 ± 1.4	[171.4, 180.6]	
ior	178.1 ± 1.1	[173.3, 180.6]	0.038
or	177.9 ± 1.1	[173.9, 180.9]	

Table 1. Mean CSJ angle per quadrant

Figure 2. Individual CSJ angle in each sector in all 105 eyes (colour lines). Corresponding mean CSJ angle (black line) and error bars (in light gray) indicating ± standard error are also shown.